Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Placental Pathology Associated with Household Air Pollution in a Cohort of Pregnant Women from Dar es Salaam, Tanzania

Blair J. Wylie, Emmanuel Matechi, Yahya Kishashu, Wafaie Fawzi, Zul Premji, Brent A. Coull, Russ Hauser, Majid Ezzati, and Drucilla Roberts

Table of Contents

Details of carbon monoxide exposure measurement

Details of fine particulate matter exposure measurement

Figure S1: Identification of the study cohort. Of the 239 subjects enrolled into the primary study on household air pollution among pregnant women in Dar es Salaam, Tanzania, 123 were excluded secondary to a lack of available placental slides. Of the remaining 116, all had available CO measurements. Only 79 of the 116 subjects had available and valid $PM_{2.5}$ measurements. The reasons for excluding $PM_{2.5}$ measurements from the other 37 subjects included: 1 inadequate sampling time, 1 misplaced filters, 4 pump errors, and 31 with implausible weights secondary to incorrect pump setup by one field worker.

Table S1: Placental pathology by particulate matter exposure, unadjusted and adjusted models

Table S2: Placental pathology by carbon monoxide exposure, unadjusted and adjusted models

References